

ORIGINAL ARTICLE

Knowledge, Attitude and Practice Pattern regarding Diabetic Retinopathy among the patients of Diabetes Mellitus presenting to a Tertiary Health Care Center in Uttarakhand

Renu Dhasmana¹, Ruchi Juyal², Hennaav Kaur Dhillon³

¹Professor, Department of Ophthalmology, Himalayan Institute of Medical Sciences, Swami Rama Himalayan University, Swami Ram Nagar, Jolly Grant, Dehradun; ²Professor, Department of Community Medicine, Himalayan Institute of Medical Sciences, Swami Rama Himalayan University, Swami Ram Nagar, Jolly Grant, Dehradun; ³Senior Resident, Department of Ophthalmology, Himalayan Institute of Medical Sciences, Swami Rama Himalayan University, Swami Ram Nagar, Jolly Grant, Dehradun

Abstract	Introduction	Methodology	Results	Conclusion	References	Citation	Tables / Figures
--------------------------	------------------------------	-----------------------------	-------------------------	----------------------------	----------------------------	--------------------------	----------------------------------

Corresponding Author

Dr Hennaav Kaur Dhillon, MS, Senior Resident, Department of Ophthalmology, Himalayan Institute of Medical Sciences, Swami Rama Himalayan University, Swami Ram Nagar, Jolly Grant, Dehradun, Uttarakhand.

E Mail ID: hennaavdhillon@gmail.com



Citation

Dhasmana R, Juyal R, Dhillon HK. Knowledge, attitude and practice pattern regarding diabetic retinopathy among the patients of diabetes mellitus presenting to a tertiary health care center in Uttarakhand. Indian J Comm Health. 2022;34(1):30-35. <https://doi.org/10.47203/IJCH.2022.v34i01.007>

Source of Funding: Nil Conflict of Interest: None declared

Article Cycle

Received: 16/10/2021; Revision: 21/01/2022; Accepted: 03/03/2022; Published: 31/03/2022

This work is licensed under a [Creative Commons Attribution 4.0 International License](#). ©The Author(s). 2022 Open Access

Abstract

Introduction: Diabetes Mellitus (DM) is an established silent killer in the developed world and is slowly increasing the burden of morbidity in the developing nations as well. Diabetic retinopathy (DR) screening protocols are not standardized in the developing world and not available to many practitioners. **Methods:** This is an observational analytical study aimed to assess the levels of knowledge, attitude and practice patterns regarding DR among the patients of DM in Uttarakhand. These were correlated with the educational status, socio-economic status, duration of DM, residence and gender. **Results:** Higher levels of knowledge which were statistically significant were observed as the socioeconomic status rose, with the increase in the duration of DM and with higher levels of educational status. A corresponding positive tone was also seen with respect to attitude and practice. However, the levels of knowledge did not correspond to the levels of practice. **Conclusion:** Integration of the health services along with the educational system would be an advancing step in spreading awareness about the disease entity itself and its morbid consequences. This would create an environment of high awareness and screening protocols will be easily streamlined.

Keywords

Diabetic Retinopathy; Diabetes Mellitus; Knowledge; Awareness; Attitude; Practice

Introduction

Diabetes Mellitus (DM), particularly type II, is a major public health concern worldwide. There will be an alarming increase in the population with type II DM both in the developed as well as the developing world as predicted by the World Health Organisation (WHO) (1). By the year 2030, the figures are estimated to increase by approximately 46% in the developed world and at a whopping 150% in the developing world (2). Being a lifelong disease, it reduces the quality of life and increases the morbidity and mortality. At the same time, the disease and its complications pose a huge economic burden for patients and their family and the society as a whole (3).

The report of the WHO states that India harbours the maximum number of diabetic patients in the world. Blindness and visual impairment due to Diabetic Retinopathy (DR) is preventable if detected early(1). DR screening protocols in most places are either not available, or not strictly adhered to.

Aims & Objectives

1. To determine the general awareness of DR among the patients of DM
2. To correlate these levels of awareness with the socio-economic and educational status and the duration of diabetes.

Material & Methods

Study Type: It was an observational analytical and non-interventional study.

Study Population and consent: A total of 200 diagnosed diabetes patients were included in the study after obtaining a written informed consent.

Study Duration: This study was conducted over a period of 12 months.

Inclusion criteria: All the patients above 10 years of age having diabetes mellitus

Exclusion criteria: Patients with corneal opacities and cataract in which the fundus was not visible.

Study Protocol: After carrying out a detailed search of literature on guidelines for conducting a Knowledge, Attitude and Practice (KAP) study and KAP questionnaire was prepared in English to suit the target population. A detailed interview was conducted with the patient regarding the patient's diabetic status, lifestyle habits and awareness of patient regarding DR in a pre-set format. Assessment of the patient included best corrected visual acuity (BCVA), slit lamp examination for anterior chamber and iris neovascularisation. Examination of the fundus was carried out with +90D slit lamp bio microscopy.

The study subjects were divided in rural and urban areas according to the criteria mentioned in the census 2011 (4). The study subjects were divided into following classes (Highly skilled, skilled, semi-skilled, un-skilled, housewives) based on their occupation as per the Minimum Wages Act 1948 (5). Highly skilled and skilled were clubbed together and the house wives were taken as a separate entity. Patients were also sub divided into subgroups accordingly to their educational status (Illiterate, literate up to class 8, literate up to class 12th, and graduates and above) as defined by UNESCO (2003) (6). Patients were also divided according to their socioeconomic status (poor, lower middle, middle, upper middle and rich) based on their monthly family income by McKinsey (2005) (7).

The level of knowledge about DR among the patients was assessed according the occupation, education, residential area, duration of DM and the economic status. Five questions were utilised:

1. Can controlled diabetic have eye problems?
2. Is DR a common health problem?
3. Can DR be prevented?
4. Is DR curable?
5. Can DR be prevented with early treatment of DM?

The attitude of the diabetic patients was assessed on the need of annual ocular examination and the response was graded as yes, no and don't know.

Ethical approval: The study adhered to the Declaration of Helsinki and former approval from the institutional ethical committee was taken.

Data Management and Statistical analysis: Interpretation and analysis of the data obtained was carried out using

SPSS version 19; SPSS Inc Chicago, Illinois, U.S.A. Chi square tests were applied to evaluate relation between awareness and unawareness among study subjects. A "P" value of <0.05 was taken to be significant.

Results

Correlation of levels of knowledge with various parameters: As the duration of diabetes increased, the knowledge about controlled diabetics having eye problems increased from 7 (10.6%) >5 years to 7 (36.8%) of 16-20 years. Knowledge about the curability of DR increased from 16 (24.2%) in >5 years to 11 (57.8%) in 16-20 years of duration of DM. The knowledge regarding the prevention of DR by timely intervention of DM was, 14 (21.2%) among the >5 years to 13 (31.7%) among 11 – 15 years and 4 (21.1%) in 16-20 years.

The level of knowledge about DR according to occupation, educational status and socioeconomic status is depicted in (Table 1), (Table 2) and (Table 3) respectively. The patients were also asked if they felt an annual ocular examination was necessary in the patients of DM. Out of the 200 surveyed patients, 51 (25.5%) had knowledge about the necessity of the yearly ocular examination out of which 32 (62.7%) were males and 19 (37.3%) were females. A total of 88 (44%) patients were not aware about the necessity of ocular screening. A statistically significant difference was seen in the knowledge about the necessity of yearly ocular examination ($p=0.0172$)

Correlations of Attitude towards the disease entity with various parameters: The study group that underwent regular ocular examination was further assessed according to location, occupation and education. 41 (20.5%) population underwent regular ocular examination out of which 9 (21.9%) belonged to rural areas where as 32 (78.1%) belonged to urban areas. Population not undergoing regular ocular examination comprised a total of 98 (61.6%) belonging to rural areas and 61 (38.4%) belonged to urban areas. There as a statistical significant difference of patients undergoing regular ocular examination in rural and urban areas ($p=0.0000$). The attitude on importance of yearly ocular examination on the basis of gender is depicted in (Table 4).

Correlation of practice levels towards the disease entity with various parameters: There was a statistically significant difference seen among the patients undergoing regular ocular examination belonging to various occupational groups, with more awareness among the skilled and semi-skilled groups ($p=0.0000$) as depicted in (Figure 1). Similar statistically significant results have been observed on the basis of educational status ($p=0.009$) as depicted in (Figure 2).

The practice of diabetic patients who aware of the necessity of regular ocular examination, referring other diabetic patients for ocular check-up was also assessed and is depicted in (Table 4) and (Figure 3).

Discussion

Diabetic retinopathy originally thought as to be a disease of the developed world has slowly crept into the developing countries and is forming a major share of lifestyle diseases. It is becoming a leading cause of blindness in the adults in the productive age group. With the current developments in the healthcare system the life span of these diabetic patients is increasing and hence the burden of the ocular morbidity is increasing exponentially which will be difficult to manage with the current health care resources. It is well established that early detection is the key to prevention of this ocular morbidity. The goal is within our reach only if there is adequate awareness of the disease and appropriate health care facilities are made available. Awareness creation in the community is the key step in any program aimed to prevent DR.

In the current study it was observed that knowledge about various aspects of DR was significantly higher in the study subjects who were graduates and above (26.5%), this correlates well with other studies conducted by Thapa et al in Nepal (8), Kadri et al (9), and Addir KR et al in Malaysia (10), where highest levels of awareness about DR was seen in the higher education group.

A statistically significant correlation in the knowledge about DR was seen in people living in urban areas in comparison to the rural areas. This could be due to the fact that people living in urban areas are more educated as compared to the rural population. Similar results were seen in studies conducted in other parts of India by Rani PK et al (11), Mahesh G et al (12) and Mithal S et al (13).

In the current study a significant association was found between the duration of DM and the know of DR. Similar results were also obtained in the studies conducted in south India (8), and by Mohammed et al in Nigeria (14) and by Addoor et al in Malaysia (15). However, no correlation was found between the duration of DM and the levels of knowledge in a study conducted by Thapa et al in Nepal (8).

In the present study group, knowledge about the various aspects of DR was statistically correlated with the economic status of the patients. This result correlated well with several studies conducted in South India by Rani et al (11) and Dandona et al (16).

A statistically significant difference was seen in the level of knowledge about the treatment of DR with increase in the level of education ($p=0.035$), occupation with more among the skilled subjects ($p=0.000$) and residential area ($p=0.0005$). In the current study patients with graduate degree were more aware. A statistically insignificant level of awareness about treatment of DR was seen in our study with economic status ($p=0.546$). Similar results can be

observed in other studies conducted in India by Sihota et al (2), Mahesh et al (12), Kadri et al (9) and Singh et al (17). In the present study out of the 39.5% of patients aware of ocular involvement among diabetic patients, 20.5% underwent ocular examination. Similar results were seen in studies conducted in India by Mahesh et al (12) and by Mohammed et al in Nigeria (14) and Addoor et al in Malaysia (15). There was little or no knowledge of retinopathy risk factors or the need for early detection through screening.

In a study conducted by Rani et al in South India (11), a higher percentage (63.5%) of people aware about ocular involvement underwent regular ocular examination. Similar results were seen in studies conducted in USA by Klein et al (18) and by Tapp et al in Australia (19).

Conclusion

Integration of the health services along with the educational system would be an advancing step in spreading awareness about the disease entity itself and its morbid consequences. This would create an environment of high awareness and screening protocols will be easily streamlined. Greater involvement of paramedical staff, grass root levels health workers such as ASHA workers, ANMs and MHW at primary and community health centers would help in increasing awareness among the masses. Cooperation from the media houses both print and electronic is the need of the hour to create more awareness. In the wake of social media, campaigns promoting the timely screening of DM as well as early ocular examination in DM is a welcoming step that is much needed.

Recommendation

In a developing country such as ours where the incidence of DM is increasing by the day, it is important to raise awareness about the long-term side effects of the illness. It is also essential to create awareness regarding screening protocols at regular intervals to avoid diagnosis at vision threatening levels. In addition to being beneficial to the patient, it contributes to alleviating the financial burden over the patient, family and the society as a whole.

Relevance of the study

With the increase in the number of cases of DM and consequently DR, it is important to know the level of awareness regarding the disease and its management protocols among the general population.

Authors Contribution

RD: Conception, design and accusation of data and drafting of the manuscript and final approval to the published version. RJ: Conception, design and accusation of data and drafting of the manuscript and final approval to the published version. HKD: Conception, design and accusation of data and drafting of the manuscript and final approval to the published version

References

- Park K. Textbook of preventive and social medicine. 16th ed. Jabapur: 2000.
- Sihota R, Tandon R. Parsons’ diseases of the eye. 19th ed. Harcourt: Butterworth-heinemann 2003.
- Gupta V. Type 2 Diabetes Mellitus in India. http://sancd.org/uploads/pdf/factsheet_diabetes_Aug_10_2011. Accessed on 23/7/2020
- Ministry of Home Affairs, Government of India. <http://censuindia.gov.in/Metadata/Metada.htm>. October 1 2011. Accessed on 23/7/2020
- Labour and employment department, Tamil Nadu. www.tn.gov.in/gosdb/gorders/labour/labemp_e2d_37_2011. Accessed on 23/7/2020
- Govinds R, Biswal K. Mapping literacy in India: Who are the illiterates and where do we find them? EFA Global Monitoring Report 2006, Literacy for Life. 2005. Accessed on 23/7/2020
- McKinsey. Socio-economic status of developing India. <http://www.mckinsey.com/locations/india/mckinseyonindia/>. March 12 2005.
- Thapa R, Paudyal G, Shrestha MK, Gurung R, Ruit S. Prevalence of diabetic retinopathy following a community screening for diabetes. Nepal Medical College Journal.2008; 10 (3): 160-3.
- Kadri R. Awareness of diabetic and hypertensive eye disease in public. Int J Biol Med Res. 2011;2(2):533-5.
- Addoor KR, Bhandary SV, Khanna R. Assessment of awareness of Diabetic Retinopathy among the Diabetics attending the peripheral diabetes clinics in Melaka, Malaysia. Med J Malaysia.2011;66(1):48-52.
- Rani PK, Raman R,Subramani S, Perumal G, Kumar Manikkavel G, Sharma T. Knowledge of diabetes and diabetic retinopathy among rural population in India, and the influence of knowledge of diabetic retinopathy on attitude and practice. Rural and remote health. 2008; 8;830.
- Mahesh G, Elias A, Giridhar A. Chengamanad Diabetic Retinopathy awareness study. Kerela Journal of Ophthalmology. 2006; 18(1): 14-21.
- Mithal S, Singh S, Gupta A. Field Survey- Assessment of Disease Magnitude and Contributory Factors for Diabetic retinopathy. Community/social Ophthalmology-I. 2011; 69: 745-8.
- Klein R, Zinnman B, Gardiener R. The relationship f Diabetic Retinopathy to pre-clinical glomerulopathy lesion in type I diabetic patients. Diabetes. 2005;54(2):527-33.
- Addoor AR, Bhandari SV, Khanna R. Assessment of awareness ofDiabetic Retinopathy among the diabetics attending the peripheral diabetic clinic in Melaka, Malaysia. Med J Malaysia. 2011; 66(1):48-52.
- Dandonna L, Dandonna R, Naduvilath TJ. Population based assessment of Diabetic Retinopathy in an urban population in Southern India. Br J Ophthalmol. 1999;83:937-40.
- Singh K, Kharbanda, B. Awareness of annual ophthalmic check-up, Treatment and prognosis in patients of diabetic retinopathy. Community/Social Ophthalmology-I. 2011;69:751-3.
- Klein R, Kristine E, Knudtson MD, Gangnon RE. The Wisconsin Epidemiological study of Diabetic Retinopathy. American Academy of Ophthalmology. 2009; 116: 1937-42.
- Tapp RJ, Grad D, Zimmet PZ. Diabetes care in an Australian population frequency of screening examination for eye and foot complication of diabetes. Diabetes care 2004; 27:688-93.

Tables

TABLE 1 KNOWLEDGE ABOUT DIABETIC RETINOPATHY ACCORDING TO OCCUPATION(N=200)

		Skilled	Semi skilled	Non skilled	Unemployed	Housewives
Controlled diabetics have eye problems	Yes	45.4%	24.2%	6.6%	33.3%	16.2%
	No	30.3%	12.1%	6.6%	0.0%	12.5%
	DK	24.3%	63.7%	86.8%	66.7%	71.3%
Chi square = 19.59		P value=0.006				
DR a common health problem	Yes	72.7%	33.3%	2.2%	44.4%	16.3%
	No	3.1%	12.1%	4.4%	0.0%	10.0%
	DK	24.2%	54.6%	93.4%	55.6%	73.7%
Chi square = 56.42		P value=0.000				
Can DR be prevented	Yes	54.5%	42.4%	4.4%	44.4%	10.0%
	No	18.8%	9.1%	0.0%	0.0%	3.7%
	DK	26.7%	48.5%	95.6%	55.6%	85.3%
Chi square = 28.37		P value=0.0001				
Is DR curable	Yes	54.5%	36.4%	4.4%	44.4%	13.7%
	No	12.1%	6.1%	0.0%	0.0%	2.5%
	DK	33.4%	57.5%	95.6%	55.6%	83.8%
Chi square = 36.25		P value=0.0000				
Can DR be prevented with early treatment of DM	Yes	54.5%	42.4%	4.4%	44.4%	10.0%
	No	18.1%	9.1%	0.0%	0.0%	3.7%
	DK	27.4%	48.5%	95.6%	55.6%	86.3%
Chi square = 43.90		P value=0.0000				

TABLE 2 CORRELATION OF LEVELS OF KNOWLEDGE WITH THE EDUCATIONAL STATUS

		Illiterate	Simple Literate	Upto class 8	Upto class 12	Graduate and above
Controlled diabetics have eye problems	Yes	6.5%	20.5%	0.0%	51.4%	31.4%
	No	9.2%	2.5%	53.3%	8.5%	22.8%
	DK	84.3%	77.0%	46.7%	40.1%	45.8%
Chi square = 35.35		P value=0.003				
DR a common health problem	Yes	10.5%	20.5%	53.3%	34.2%	40.0%
	No	3.9%	2.5%	20.0%	22.8%	0.0%
	DK	85.6%	77.0%	26.7%	43.0%	60.0%
Chi square = 22.70		P value=0.00014				
Can DR be prevented	Yes	9.2%	17.9%	20.0%	28.5%	54.2%
	No	0.0%	2.5%	0.0%	20.0%	11.4%
	DK	90.8%	79.6%	80.0%	51.5%	34.4%
Chi square = 28.05		P value=0.0001				
Is DR curable	Yes	0.0%	0.0	0.0	2.8	0.0
	No	5.2%	5.1	13.3	73.2	25.7
	DK	94.8%	94.9	86.7	64.0	74.3
Chi square = 4.74		P value=0.0315				
Can DR be prevented with early treatment of DM	Yes	9.2%	17.9%	20.0%	28.5%	54.2%
	No	0.0%	2.5%	0.0%	20.0%	11.4%
	DK	90.8%	79.6%	80.0%	51.5%	34.4%
Chi square = 28.76		P value=0.0008				

TABLE 3 CORRELATION OF LEVELS OF KNOWLEDGE WITH THE SOCIO ECONOMIC STATUS

		Poor	Lower Middle	Middle	Upper Middle	Rich
Controlled diabetics have eye problems	Yes	6.5%	17.9%	13.3%	51.4%	31.4%
	No	9.2%	5.1%	40.0%	14.2%	28.5%
	DK	84.3%	77.0%	46.7%	34.4%	40.1%
Chi square = 31.53		P value=0.0002				
DR a common health problem	Yes	10.5%	20.5%	53.3%	27.1%	40.0%
	No	5.2%	2.5%	20.0%	25.8%	5.7%
	DK	84.3%	77.0%	26.7%	37.1%	54.3%
Chi square = 25.26		P value=0.0004				
Can DR be prevented	Yes	9.2%	15.3%	20.0%	28.5%	54.2%
	No	2.6%	5.1%	13.3%	20.0%	11.4%
	DK	88.2%	79.6%	66.7%	51.5%	34.4%
Chi square = 29.13		P value=0.0007				
Is DR curable	Yes	5.2%	5.1%	13.3%	11.4%	11.4%
	No	5.2%	5.1%	13.3%	28.5%	25.7%
	DK	89.6%	89.8%	73.4%	60.1%	62.9%
Chi square = 3.07		P value=0.546				
Can DR be prevented with early treatment of DM	Yes	7.8	17.9	20.0	28.5	42.8
	No	2.6	2.5	13.3	20.0	17.1
	DK	89.6	79.6	66.7	51.5	40.1
Chi square = 19.70		P value=0.0005				

TABLE 4 ATTITUDE ON IMPORTANCE OF YEARLY OCULAR EXAMINATION

Yearly Ocular Examination	Males N(%)	Females N(%)	Total N (%)	P Value
Yes	32 (67.7)	19 (37.3)	51 (25.5)	0.017
No	36 (59.9)	25 (41)	61 (30.5)	
Don't Know	42 (47.7)	46 (52.3)	88 (44.0)	

Figures

FIGURE 1 PRACTICE PATTERNS ON THE BASIS OF OCCUPATIONAL STATUS




FIGURE 2 PRACTICE PATTERNS ON THE BASIS OF EDUCATIONAL STATUS




FIGURE 3 PRACTICE OF REFERRING OTHER DIABETIC PATIENTS FOR OPHTHALMOLOGICAL EXAMINATION (N=79)

